

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00007XX
TrigonometryExercise6 for No.9562

$$No1 = \begin{bmatrix} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 3 = 0) \\ .3 = (\cos(2 \theta) = 0) & .4 = (\sin(2 \theta) = 1) \end{bmatrix}$$

$$No2 = \begin{bmatrix} .1 = (2 [\cos(\theta)]^2 - \cos(\theta) - 1 = 0) & .2 = (2 [\cos(\theta)]^2 - \sin(\theta) - 1 = 0) \\ .3 = (2 [\sin(\theta)]^2 - \sin(\theta) = 0) & .4 = (\sin(\theta) - \cos(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \sin(\theta) = \tan(\theta)) & .6 = (4 [\sin(\theta)]^3 - 3 \sin(\theta) = 0) \end{bmatrix}$$

$$No3 = \begin{bmatrix} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 1 = 0) \\ .3 = (\sin(4 \theta) = 0) & .4 = (\cos(3 \theta) = 1) \end{bmatrix}$$

$$No4 = \begin{bmatrix} .1 = (2 [\cos(\theta)]^2 + 3 \cos(\theta) + 1 = 0) & .2 = (2 [\sin(\theta)]^2 + 3 \cos(\theta) - 3 = 0) \\ .3 = (2 [\sin(\theta)]^2 + \sin(\theta) = 0) & .4 = (\cos(\theta) - \sin(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \cos(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - 3 \cos(\theta) = 0) \end{bmatrix}$$

$$No5 = \begin{bmatrix} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\sin(\theta)]^2 = 3) \\ .3 = (\tan(\theta) \cos(\theta) = \tan(\theta)) & .4 = (3 [\tan(\theta)]^2 - 1 = 0) \end{bmatrix}$$

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00008XX
TrigonometryExercise6 for No.9721

$$No1 = \begin{bmatrix} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 1 = 0) \\ .3 = (\cos(3 \theta) = 0) & .4 = (\cos(4 \theta) = 1) \end{bmatrix}$$

$$No2 = \begin{bmatrix} .1 = (2 [\sin(\theta)]^2 + 3 \sin(\theta) + 1 = 0) & .2 = (2 [\cos(\theta)]^2 + 3 \sin(\theta) - 3 = 0) \\ .3 = (2 [\sin(\theta)]^2 - \sin(\theta) = 0) & .4 = (\cos(\theta) - \sin(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \sin(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - 3 \cos(\theta) = 0) \end{bmatrix}$$

$$No3 = \begin{bmatrix} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 1 = 0) \\ .3 = (\cos(2 \theta) = 0) & .4 = (\sin(2 \theta) = 1) \end{bmatrix}$$

$$No4 = \begin{bmatrix} .1 = (2 [\sin(\theta)]^2 - \sin(\theta) - 1 = 0) & .2 = (2 [\sin(\theta)]^2 - \cos(\theta) - 1 = 0) \\ .3 = (2 [\cos(\theta)]^2 + \cos(\theta) = 0) & .4 = (\sin(\theta) - \cos(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \sin(\theta) = \sin(\theta)) & .6 = (4 [\sin(\theta)]^3 - 3 \sin(\theta) = 0) \end{bmatrix}$$

$$No5 = \begin{bmatrix} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 = 3) \\ .3 = (2 \tan(\theta) \cos(\theta) = \tan(\theta)) & .4 = ([\tan(\theta)]^2 - 1 = 0) \end{bmatrix}$$

X [Page = 0004] XXX

:
:
:
:
:
:
:
:
:
:

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00013XX
TrigonometryExercise6 for No.9854

$$\begin{aligned}
 \text{No1} &= \left[\begin{array}{ll} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 3 = 0) \\ .3 = (\cos(4\theta) = 0) & .4 = (\cos(3\theta) = 1) \end{array} \right] \\
 \text{No2} &= \left[\begin{array}{ll} .1 = (2 [\cos(\theta)]^2 + \cos(\theta) - 1 = 0) & .2 = (2 [\cos(\theta)]^2 + 3 \sin(\theta) - 3 = 0) \\ .3 = (2 [\cos(\theta)]^2 + \cos(\theta) = 0) & .4 = (\cos(\theta) - \sin(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \cos(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - 3 \cos(\theta) = 0) \end{array} \right] \\
 \text{No3} &= \left[\begin{array}{ll} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 1 = 0) \\ .3 = (\cos(2\theta) = 0) & .4 = (\sin(4\theta) = 1) \end{array} \right] \\
 \text{No4} &= \left[\begin{array}{ll} .1 = (2 [\cos(\theta)]^2 - \cos(\theta) - 1 = 0) & .2 = (2 [\sin(\theta)]^2 - 3 \cos(\theta) - 3 = 0) \\ .3 = (2 [\cos(\theta)]^2 - \cos(\theta) = 0) & .4 = (\sin(\theta) + \cos(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \cos(\theta) = \cos(\theta)) & .6 = (4 [\cos(\theta)]^3 - \cos(\theta) = 0) \end{array} \right] \\
 \text{No5} &= \left[\begin{array}{ll} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\sin(\theta)]^2 = 1) \\ .3 = (2 \tan(\theta) \sin(\theta) = \tan(\theta)) & .4 = ([\tan(\theta)]^2 - 1 = 0) \end{array} \right]
 \end{aligned}$$

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00014XX
TrigonometryExercise6 for No.9862

$$\begin{aligned}
 \text{No1} &= \left[\begin{array}{ll} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 3 = 0) \\ .3 = (\sin(2\theta) = 0) & .4 = (\cos(3\theta) = 1) \end{array} \right] \\
 \text{No2} &= \left[\begin{array}{ll} .1 = (2 [\sin(\theta)]^2 + 3 \sin(\theta) + 1 = 0) & .2 = (2 [\cos(\theta)]^2 - \sin(\theta) - 1 = 0) \\ .3 = (2 [\sin(\theta)]^2 + \sin(\theta) = 0) & .4 = (\sin(\theta) - \cos(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \cos(\theta) = \cos(\theta)) & .6 = (4 [\cos(\theta)]^3 - \cos(\theta) = 0) \end{array} \right] \\
 \text{No3} &= \left[\begin{array}{ll} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 3 = 0) \\ .3 = (\cos(2\theta) = 0) & .4 = (\cos(4\theta) = 1) \end{array} \right] \\
 \text{No4} &= \left[\begin{array}{ll} .1 = (2 [\cos(\theta)]^2 - 3 \cos(\theta) + 1 = 0) & .2 = (2 [\sin(\theta)]^2 + \cos(\theta) - 1 = 0) \\ .3 = (2 [\sin(\theta)]^2 - \sin(\theta) = 0) & .4 = (\cos(\theta) - \sin(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \sin(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - 3 \cos(\theta) = 0) \end{array} \right] \\
 \text{No5} &= \left[\begin{array}{ll} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 = 1) \\ .3 = (\tan(\theta) \sin(\theta) = \sin(\theta)) & .4 = ([\tan(\theta)]^2 - 1 = 0) \end{array} \right]
 \end{aligned}$$

X [Page = 0007] XXX

:
:
:
:
:
:
:
:
:
:
:

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00017XX

TrigonometryExercise6 for No.10532

$$No1 = \begin{bmatrix} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 1 = 0) \\ .3 = (\cos(2 \theta) = 0) & .4 = (\sin(2 \theta) = 1) \end{bmatrix}$$

$$No2 = \begin{bmatrix} .1 = (2 [\sin(\theta)]^2 + \sin(\theta) - 1 = 0) & .2 = (2 [\cos(\theta)]^2 - 3 \sin(\theta) - 3 = 0) \\ .3 = (2 [\cos(\theta)]^2 + \cos(\theta) = 0) & .4 = (\cos(\theta) - \sin(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \sin(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - 3 \cos(\theta) = 0) \end{bmatrix}$$

$$No3 = \begin{bmatrix} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 3 = 0) \\ .3 = (\sin(3 \theta) = 0) & .4 = (\sin(4 \theta) = 1) \end{bmatrix}$$

$$No4 = \begin{bmatrix} .1 = (2 [\sin(\theta)]^2 - 3 \sin(\theta) + 1 = 0) & .2 = (2 [\sin(\theta)]^2 + \cos(\theta) - 1 = 0) \\ .3 = (2 [\sin(\theta)]^2 - \sin(\theta) = 0) & .4 = (\sin(\theta) - \cos(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \sin(\theta) = \tan(\theta)) & .6 = (4 [\sin(\theta)]^3 - 3 \sin(\theta) = 0) \end{bmatrix}$$

$$No5 = \begin{bmatrix} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 = 3) \\ .3 = (\tan(\theta) \cos(\theta) = \tan(\theta)) & .4 = ([\tan(\theta)]^2 - 3 = 0) \end{bmatrix}$$

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00018XX

TrigonometryExercise6 for No.10688

$$No1 = \begin{bmatrix} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 3 = 0) \\ .3 = (\sin(3 \theta) = 0) & .4 = (\cos(3 \theta) = 1) \end{bmatrix}$$

$$No2 = \begin{bmatrix} .1 = (2 [\cos(\theta)]^2 + 3 \cos(\theta) + 1 = 0) & .2 = (2 [\cos(\theta)]^2 + \sin(\theta) - 1 = 0) \\ .3 = (2 [\cos(\theta)]^2 + \cos(\theta) = 0) & .4 = (\sin(\theta) + \cos(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \sin(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - \cos(\theta) = 0) \end{bmatrix}$$

$$No3 = \begin{bmatrix} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 1 = 0) \\ .3 = (\sin(2 \theta) = 0) & .4 = (\sin(4 \theta) = 1) \end{bmatrix}$$

$$No4 = \begin{bmatrix} .1 = (2 [\cos(\theta)]^2 - 3 \cos(\theta) + 1 = 0) & .2 = (2 [\sin(\theta)]^2 - \cos(\theta) - 1 = 0) \\ .3 = (2 [\cos(\theta)]^2 - \cos(\theta) = 0) & .4 = (\sin(\theta) - \cos(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \cos(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - 3 \cos(\theta) = 0) \end{bmatrix}$$

$$No5 = \begin{bmatrix} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\sin(\theta)]^2 = 1) \\ .3 = (\tan(\theta) \cos(\theta) = \tan(\theta)) & .4 = ([\tan(\theta)]^2 - 1 = 0) \end{bmatrix}$$

X [Page = 0009] XXX

:
:
:
:
:
:
:
:
:
:
:

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00027XX
TrigonometryExercise6 for No.12278

$$No1 = \left[\begin{array}{ll} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 1 = 0) \\ .3 = (\sin(2\theta) = 0) & .4 = (\sin(3\theta) = 1) \end{array} \right]$$

$$No2 = \left[\begin{array}{ll} .1 = (2 [\cos(\theta)]^2 - 3 \cos(\theta) + 1 = 0) & .2 = (2 [\sin(\theta)]^2 - \cos(\theta) - 1 = 0) \\ .3 = (2 [\cos(\theta)]^2 + \cos(\theta) = 0) & .4 = (\sin(\theta) + \cos(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \cos(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - \cos(\theta) = 0) \end{array} \right]$$

$$No3 = \left[\begin{array}{ll} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 1 = 0) \\ .3 = (\cos(2\theta) = 0) & .4 = (\sin(4\theta) = 1) \end{array} \right]$$

$$No4 = \left[\begin{array}{ll} .1 = (2 [\cos(\theta)]^2 - \cos(\theta) - 1 = 0) & .2 = (2 [\cos(\theta)]^2 - 3 \sin(\theta) - 3 = 0) \\ .3 = (2 [\sin(\theta)]^2 - \sin(\theta) = 0) & .4 = (\cos(\theta) - \sin(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \sin(\theta) = \sin(\theta)) & .6 = (4 [\sin(\theta)]^3 - \sin(\theta) = 0) \end{array} \right]$$

$$No5 = \left[\begin{array}{ll} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 = 3) \\ .3 = (\tan(\theta) \cos(\theta) = \cos(\theta)) & .4 = ([\tan(\theta)]^2 - 3 = 0) \end{array} \right]$$

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00028XX
TrigonometryExercise6 for No.12482

$$No1 = \left[\begin{array}{ll} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 1 = 0) \\ .3 = (\cos(2\theta) = 0) & .4 = (\sin(2\theta) = 1) \end{array} \right]$$

$$No2 = \left[\begin{array}{ll} .1 = (2 [\cos(\theta)]^2 - \cos(\theta) - 1 = 0) & .2 = (2 [\sin(\theta)]^2 + 3 \cos(\theta) - 3 = 0) \\ .3 = (2 [\sin(\theta)]^2 - \sin(\theta) = 0) & .4 = (\cos(\theta) - \sin(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \sin(\theta) = \sin(\theta)) & .6 = (4 [\sin(\theta)]^3 - 3 \sin(\theta) = 0) \end{array} \right]$$

$$No3 = \left[\begin{array}{ll} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 3 = 0) \\ .3 = (\cos(4\theta) = 0) & .4 = (\cos(3\theta) = 1) \end{array} \right]$$

$$No4 = \left[\begin{array}{ll} .1 = (2 [\sin(\theta)]^2 + 3 \sin(\theta) + 1 = 0) & .2 = (2 [\cos(\theta)]^2 - \sin(\theta) - 1 = 0) \\ .3 = (2 [\sin(\theta)]^2 + \sin(\theta) = 0) & .4 = (\sin(\theta) + \cos(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \cos(\theta) = \cos(\theta)) & .6 = (4 [\cos(\theta)]^3 - \cos(\theta) = 0) \end{array} \right]$$

$$No5 = \left[\begin{array}{ll} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 = 1) \\ .3 = (\tan(\theta) \sin(\theta) = \tan(\theta)) & .4 = (3 [\tan(\theta)]^2 - 1 = 0) \end{array} \right]$$

X [Page = 0014] XXX

:
:
:
:
:
:
:
:
:
:
:
:
:

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00037XX

TrigonometryExercise6 for No.645104

$$No1 = \left[\begin{array}{ll} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 3 = 0) \\ .3 = (\cos(3 \theta) = 0) & .4 = (\cos(4 \theta) = 1) \end{array} \right]$$

$$No2 = \left[\begin{array}{ll} .1 = (2 [\cos(\theta)]^2 - \cos(\theta) - 1 = 0) & .2 = (2 [\sin(\theta)]^2 + 3 \cos(\theta) - 3 = 0) \\ .3 = (2 [\cos(\theta)]^2 - \cos(\theta) = 0) & .4 = (\sin(\theta) - \cos(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \sin(\theta) = \tan(\theta)) & .6 = (4 [\sin(\theta)]^3 - 3 \sin(\theta) = 0) \end{array} \right]$$

$$No3 = \left[\begin{array}{ll} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 3 = 0) \\ .3 = (\sin(2 \theta) = 0) & .4 = (\sin(4 \theta) = 1) \end{array} \right]$$

$$No4 = \left[\begin{array}{ll} .1 = (2 [\cos(\theta)]^2 + 3 \cos(\theta) + 1 = 0) & .2 = (2 [\cos(\theta)]^2 - \sin(\theta) - 1 = 0) \\ .3 = (2 [\cos(\theta)]^2 + \cos(\theta) = 0) & .4 = (\cos(\theta) - \sin(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \sin(\theta) = \sin(\theta)) & .6 = (4 [\cos(\theta)]^3 - 3 \cos(\theta) = 0) \end{array} \right]$$

$$No5 = \left[\begin{array}{ll} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\cos(\theta)]^2 = 1) \\ .3 = (\tan(\theta) \cos(\theta) = \cos(\theta)) & .4 = (3 [\tan(\theta)]^2 - 1 = 0) \end{array} \right]$$

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00038XX

TrigonometryExercise6 for No.645105

$$No1 = \left[\begin{array}{ll} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 1 = 0) \\ .3 = (\sin(4 \theta) = 0) & .4 = (\cos(4 \theta) = 1) \end{array} \right]$$

$$No2 = \left[\begin{array}{ll} .1 = (2 [\sin(\theta)]^2 - 3 \sin(\theta) + 1 = 0) & .2 = (2 [\cos(\theta)]^2 - \sin(\theta) - 1 = 0) \\ .3 = (2 [\sin(\theta)]^2 - \sin(\theta) = 0) & .4 = (\sin(\theta) + \cos(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \cos(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - \cos(\theta) = 0) \end{array} \right]$$

$$No3 = \left[\begin{array}{ll} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 1 = 0) \\ .3 = (\cos(3 \theta) = 0) & .4 = (\cos(2 \theta) = 1) \end{array} \right]$$

$$No4 = \left[\begin{array}{ll} .1 = (2 [\sin(\theta)]^2 + 3 \sin(\theta) + 1 = 0) & .2 = (2 [\sin(\theta)]^2 + \cos(\theta) - 1 = 0) \\ .3 = (2 [\cos(\theta)]^2 + \cos(\theta) = 0) & .4 = (\sin(\theta) - \cos(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \cos(\theta) = \tan(\theta)) & .6 = (4 [\sin(\theta)]^3 - 3 \sin(\theta) = 0) \end{array} \right]$$

$$No5 = \left[\begin{array}{ll} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 = 3) \\ .3 = (\tan(\theta) \sin(\theta) = \sin(\theta)) & .4 = (3 [\tan(\theta)]^2 - 1 = 0) \end{array} \right]$$

X [Page = 0019] XXX

:
:
:
:
:
:
:
:
:
:
:
:

XXX
:
:
:
:
:
X Math@MUT XX
TrigonometryExercise6 for No.645108

$$No1 = \left[\begin{array}{l} .1 = (2 \sin(\theta) - 1 = 0) \quad .2 = (4 [\cos(\theta)]^2 - 3 = 0) \\ .3 = (\cos(3\theta) = 0) \quad .4 = (\sin(4\theta) = 1) \end{array} \right]$$

$$No2 = \left[\begin{array}{l} .1 = (2 [\sin(\theta)]^2 - \sin(\theta) - 1 = 0) \quad .2 = (2 [\cos(\theta)]^2 - \sin(\theta) - 1 = 0) \\ .3 = (2 [\cos(\theta)]^2 - \cos(\theta) = 0) \quad .4 = (\sin(\theta) + \cos(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \cos(\theta) = \tan(\theta)) \quad .6 = (4 [\cos(\theta)]^3 - \cos(\theta) = 0) \end{array} \right]$$

$$No3 = \left[\begin{array}{l} .1 = (2 \sin(\theta) + 1 = 0) \quad .2 = (4 [\sin(\theta)]^2 - 1 = 0) \\ .3 = (\sin(3\theta) = 0) \quad .4 = (\cos(2\theta) = 1) \end{array} \right]$$

$$No4 = \left[\begin{array}{l} .1 = (2 [\sin(\theta)]^2 + 3 \sin(\theta) + 1 = 0) \quad .2 = (2 [\sin(\theta)]^2 + 3 \cos(\theta) - 3 = 0) \\ .3 = (2 [\cos(\theta)]^2 + \cos(\theta) = 0) \quad .4 = (\sin(\theta) - \cos(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \sin(\theta) = \tan(\theta)) \quad .6 = (4 [\sin(\theta)]^3 - 3 \sin(\theta) = 0) \end{array} \right]$$

$$No5 = \left[\begin{array}{l} .1 = (2 \cos(\theta) - 1 = 0) \quad .2 = (4 [\cos(\theta)]^2 = 1) \\ .3 = (\tan(\theta) \sin(\theta) = \sin(\theta)) \quad .4 = (3 [\tan(\theta)]^2 - 1 = 0) \end{array} \right]$$

XXX
:
:
:
:
:
X Math@MUT XX
TrigonometryExercise6 for No.645109

$$No1 = \left[\begin{array}{l} .1 = (2 \sin(\theta) + 1 = 0) \quad .2 = (4 [\cos(\theta)]^2 - 1 = 0) \\ .3 = (\sin(3\theta) = 0) \quad .4 = (\sin(4\theta) = 1) \end{array} \right]$$

$$No2 = \left[\begin{array}{l} .1 = (2 [\sin(\theta)]^2 + \sin(\theta) - 1 = 0) \quad .2 = (2 [\cos(\theta)]^2 - 3 \sin(\theta) - 3 = 0) \\ .3 = (2 [\sin(\theta)]^2 - \sin(\theta) = 0) \quad .4 = (\cos(\theta) - \sin(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \sin(\theta) = \tan(\theta)) \quad .6 = (4 [\sin(\theta)]^3 - \sin(\theta) = 0) \end{array} \right]$$

$$No3 = \left[\begin{array}{l} .1 = (2 \cos(\theta) - 1 = 0) \quad .2 = (4 [\cos(\theta)]^2 - 3 = 0) \\ .3 = (\cos(4\theta) = 0) \quad .4 = (\cos(2\theta) = 1) \end{array} \right]$$

$$No4 = \left[\begin{array}{l} .1 = (2 [\sin(\theta)]^2 - \sin(\theta) - 1 = 0) \quad .2 = (2 [\sin(\theta)]^2 + 3 \cos(\theta) - 3 = 0) \\ .3 = (2 [\cos(\theta)]^2 + \cos(\theta) = 0) \quad .4 = (\sin(\theta) - \cos(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \sin(\theta) = \sin(\theta)) \quad .6 = (4 [\cos(\theta)]^3 - 3 \cos(\theta) = 0) \end{array} \right]$$

$$No5 = \left[\begin{array}{l} .1 = (2 \sin(\theta) - 1 = 0) \quad .2 = (4 [\sin(\theta)]^2 = 1) \\ .3 = (\tan(\theta) \sin(\theta) = \tan(\theta)) \quad .4 = ([\tan(\theta)]^2 - 3 = 0) \end{array} \right]$$

X [Page = 0021] XX
:
:
:
:
:
:
:
:
:
:
:

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00049XX

TrigonometryExercise6 for No.645116

$$No1 = \begin{bmatrix} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 1 = 0) \\ .3 = (\sin(4 \theta) = 0) & .4 = (\sin(2 \theta) = 1) \end{bmatrix}$$

$$No2 = \begin{bmatrix} .1 = (2 [\cos(\theta)]^2 + \cos(\theta) - 1 = 0) & .2 = (2 [\cos(\theta)]^2 + \sin(\theta) - 1 = 0) \\ .3 = (2 [\sin(\theta)]^2 + \sin(\theta) = 0) & .4 = (\sin(\theta) + \cos(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \sin(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - 3 \cos(\theta) = 0) \end{bmatrix}$$

$$No3 = \begin{bmatrix} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 3 = 0) \\ .3 = (\cos(2 \theta) = 0) & .4 = (\cos(4 \theta) = 1) \end{bmatrix}$$

$$No4 = \begin{bmatrix} .1 = (2 [\sin(\theta)]^2 - 3 \sin(\theta) + 1 = 0) & .2 = (2 [\sin(\theta)]^2 - 3 \cos(\theta) - 3 = 0) \\ .3 = (2 [\cos(\theta)]^2 - \cos(\theta) = 0) & .4 = (\cos(\theta) - \sin(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \cos(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - \cos(\theta) = 0) \end{bmatrix}$$

$$No5 = \begin{bmatrix} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\sin(\theta)]^2 = 1) \\ .3 = (\tan(\theta) \sin(\theta) = \sin(\theta)) & .4 = ([\tan(\theta)]^2 - 3 = 0) \end{bmatrix}$$

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00050XX

TrigonometryExercise6 for No.645117

$$No1 = \begin{bmatrix} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 3 = 0) \\ .3 = (\cos(3 \theta) = 0) & .4 = (\sin(4 \theta) = 1) \end{bmatrix}$$

$$No2 = \begin{bmatrix} .1 = (2 [\cos(\theta)]^2 - 3 \cos(\theta) + 1 = 0) & .2 = (2 [\cos(\theta)]^2 + \sin(\theta) - 1 = 0) \\ .3 = (2 [\cos(\theta)]^2 + \cos(\theta) = 0) & .4 = (\sin(\theta) + \cos(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \cos(\theta) = \cos(\theta)) & .6 = (4 [\sin(\theta)]^3 - 3 \sin(\theta) = 0) \end{bmatrix}$$

$$No3 = \begin{bmatrix} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 3 = 0) \\ .3 = (\sin(2 \theta) = 0) & .4 = (\sin(3 \theta) = 1) \end{bmatrix}$$

$$No4 = \begin{bmatrix} .1 = (2 [\cos(\theta)]^2 + 3 \cos(\theta) + 1 = 0) & .2 = (2 [\cos(\theta)]^2 - \sin(\theta) - 1 = 0) \\ .3 = (2 [\cos(\theta)]^2 - \cos(\theta) = 0) & .4 = (\sin(\theta) - \cos(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \cos(\theta) = \tan(\theta)) & .6 = (4 [\sin(\theta)]^3 - \sin(\theta) = 0) \end{bmatrix}$$

$$No5 = \begin{bmatrix} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 = 1) \\ .3 = (2 \tan(\theta) \sin(\theta) = \tan(\theta)) & .4 = (3 [\tan(\theta)]^2 - 1 = 0) \end{bmatrix}$$

X [Page = 0025] XXX

:
:
:
:
:
:
:
:
:
:
:
:

XX
 :
 :
 :
 :
 X Math@MUT XXXM5/1-6400306-00051XX
 TrigonometryExercise6 for No.645118

$$No1 = \left[\begin{array}{ll} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 1 = 0) \\ .3 = (\cos(3 \theta) = 0) & .4 = (\cos(4 \theta) = 1) \end{array} \right]$$

$$No2 = \left[\begin{array}{ll} .1 = (2 [\cos(\theta)]^2 - 3 \cos(\theta) + 1 = 0) & .2 = (2 [\sin(\theta)]^2 - 3 \cos(\theta) - 3 = 0) \\ .3 = (2 [\cos(\theta)]^2 + \cos(\theta) = 0) & .4 = (\sin(\theta) + \cos(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \sin(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - \cos(\theta) = 0) \end{array} \right]$$

$$No3 = \left[\begin{array}{ll} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 1 = 0) \\ .3 = (\cos(2 \theta) = 0) & .4 = (\sin(2 \theta) = 1) \end{array} \right]$$

$$No4 = \left[\begin{array}{ll} .1 = (2 [\cos(\theta)]^2 - \cos(\theta) - 1 = 0) & .2 = (2 [\cos(\theta)]^2 - \sin(\theta) - 1 = 0) \\ .3 = (2 [\sin(\theta)]^2 - \sin(\theta) = 0) & .4 = (\cos(\theta) - \sin(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \sin(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - 3 \cos(\theta) = 0) \end{array} \right]$$

$$No5 = \left[\begin{array}{ll} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\sin(\theta)]^2 = 3) \\ .3 = (\tan(\theta) \cos(\theta) = \cos(\theta)) & .4 = (3 [\tan(\theta)]^2 - 1 = 0) \end{array} \right]$$

XX
 :
 :
 :
 :
 X Math@MUT XXXM5/1-6400306-00052XX
 TrigonometryExercise6 for No.645119

$$No1 = \left[\begin{array}{ll} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 3 = 0) \\ .3 = (\cos(4 \theta) = 0) & .4 = (\sin(3 \theta) = 1) \end{array} \right]$$

$$No2 = \left[\begin{array}{ll} .1 = (2 [\sin(\theta)]^2 - 3 \sin(\theta) + 1 = 0) & .2 = (2 [\sin(\theta)]^2 - \cos(\theta) - 1 = 0) \\ .3 = (2 [\sin(\theta)]^2 + \sin(\theta) = 0) & .4 = (\sin(\theta) + \cos(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \cos(\theta) = \tan(\theta)) & .6 = (4 [\sin(\theta)]^3 - 3 \sin(\theta) = 0) \end{array} \right]$$

$$No3 = \left[\begin{array}{ll} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 3 = 0) \\ .3 = (\sin(4 \theta) = 0) & .4 = (\sin(2 \theta) = 1) \end{array} \right]$$

$$No4 = \left[\begin{array}{ll} .1 = (2 [\cos(\theta)]^2 - \cos(\theta) - 1 = 0) & .2 = (2 [\sin(\theta)]^2 - 3 \cos(\theta) - 3 = 0) \\ .3 = (2 [\cos(\theta)]^2 - \cos(\theta) = 0) & .4 = (\sin(\theta) - \cos(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \cos(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - 3 \cos(\theta) = 0) \end{array} \right]$$

$$No5 = \left[\begin{array}{ll} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 = 1) \\ .3 = (\tan(\theta) \sin(\theta) = \sin(\theta)) & .4 = ([\tan(\theta)]^2 - 3 = 0) \end{array} \right]$$

X [Page = 0026] XXX
 :
 :
 :
 :
 :
 :
 :
 :

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00059XX

TrigonometryExercise6 for No.645126

$$\begin{aligned}
 No1 &= \left[\begin{array}{ll} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 3 = 0) \\ .3 = (\sin(3 \theta) = 0) & .4 = (\cos(3 \theta) = 1) \end{array} \right] \\
 No2 &= \left[\begin{array}{ll} .1 = (2 [\cos(\theta)]^2 - 3 \cos(\theta) + 1 = 0) & .2 = (2 [\sin(\theta)]^2 - \cos(\theta) - 1 = 0) \\ .3 = (2 [\cos(\theta)]^2 - \cos(\theta) = 0) & .4 = (\sin(\theta) - \cos(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \cos(\theta) = \tan(\theta)) & .6 = (4 [\sin(\theta)]^3 - \sin(\theta) = 0) \end{array} \right] \\
 No3 &= \left[\begin{array}{ll} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 1 = 0) \\ .3 = (\cos(4 \theta) = 0) & .4 = (\sin(2 \theta) = 1) \end{array} \right] \\
 No4 &= \left[\begin{array}{ll} .1 = (2 [\cos(\theta)]^2 + 3 \cos(\theta) + 1 = 0) & .2 = (2 [\sin(\theta)]^2 + \cos(\theta) - 1 = 0) \\ .3 = (2 [\cos(\theta)]^2 + \cos(\theta) = 0) & .4 = (\cos(\theta) - \sin(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \cos(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - 3 \cos(\theta) = 0) \end{array} \right] \\
 No5 &= \left[\begin{array}{ll} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 = 3) \\ .3 = (\tan(\theta) \sin(\theta) = \sin(\theta)) & .4 = (3 [\tan(\theta)]^2 - 1 = 0) \end{array} \right]
 \end{aligned}$$

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00060XX

TrigonometryExercise6 for No.645127

$$\begin{aligned}
 No1 &= \left[\begin{array}{ll} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 1 = 0) \\ .3 = (\cos(3 \theta) = 0) & .4 = (\cos(4 \theta) = 1) \end{array} \right] \\
 No2 &= \left[\begin{array}{ll} .1 = (2 [\sin(\theta)]^2 + 3 \sin(\theta) + 1 = 0) & .2 = (2 [\cos(\theta)]^2 + \sin(\theta) - 1 = 0) \\ .3 = (2 [\sin(\theta)]^2 + \sin(\theta) = 0) & .4 = (\sin(\theta) - \cos(\theta) = \sqrt{2}) \\ .5 = (2 \tan(\theta) \cos(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - 3 \cos(\theta) = 0) \end{array} \right] \\
 No3 &= \left[\begin{array}{ll} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\sin(\theta)]^2 - 3 = 0) \\ .3 = (\sin(2 \theta) = 0) & .4 = (\cos(2 \theta) = 1) \end{array} \right] \\
 No4 &= \left[\begin{array}{ll} .1 = (2 [\sin(\theta)]^2 + \sin(\theta) - 1 = 0) & .2 = (2 [\cos(\theta)]^2 + 3 \sin(\theta) - 3 = 0) \\ .3 = (2 [\sin(\theta)]^2 - \sin(\theta) = 0) & .4 = (\cos(\theta) - \sin(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \cos(\theta) = \tan(\theta)) & .6 = (4 [\cos(\theta)]^3 - \cos(\theta) = 0) \end{array} \right] \\
 No5 &= \left[\begin{array}{ll} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 = 3) \\ .3 = (\tan(\theta) \sin(\theta) = \sin(\theta)) & .4 = (3 [\tan(\theta)]^2 - 1 = 0) \end{array} \right]
 \end{aligned}$$

X [Page = 0030] XXX

:
:
:
:
:
:
:
:
:
:
:
:

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00061XX

TrigonometryExercise6 for No.645128

$$\begin{aligned}
No1 &= \left[\begin{array}{l} .1 = (2 \sin(\theta) + 1 = 0) \quad .2 = (4 [\cos(\theta)]^2 - 3 = 0) \\ .3 = (\sin(3\theta) = 0) \quad .4 = (\sin(2\theta) = 1) \end{array} \right] \\
No2 &= \left[\begin{array}{l} .1 = (2 [\sin(\theta)]^2 - \sin(\theta) - 1 = 0) \quad .2 = (2 [\cos(\theta)]^2 - \sin(\theta) - 1 = 0) \\ .3 = (2 [\cos(\theta)]^2 - \cos(\theta) = 0) \quad .4 = (\sin(\theta) + \cos(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \sin(\theta) = \sin(\theta)) \quad .6 = (4 [\cos(\theta)]^3 - 3 \cos(\theta) = 0) \end{array} \right] \\
No3 &= \left[\begin{array}{l} .1 = (2 \sin(\theta) - 1 = 0) \quad .2 = (4 [\sin(\theta)]^2 - 3 = 0) \\ .3 = (\cos(4\theta) = 0) \quad .4 = (\cos(2\theta) = 1) \end{array} \right] \\
No4 &= \left[\begin{array}{l} .1 = (2 [\sin(\theta)]^2 + 3 \sin(\theta) + 1 = 0) \quad .2 = (2 [\sin(\theta)]^2 + 3 \cos(\theta) - 3 = 0) \\ .3 = (2 [\cos(\theta)]^2 + \cos(\theta) = 0) \quad .4 = (\sin(\theta) - \cos(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \cos(\theta) = \tan(\theta)) \quad .6 = (4 [\sin(\theta)]^3 - \sin(\theta) = 0) \end{array} \right] \\
No5 &= \left[\begin{array}{l} .1 = (2 \cos(\theta) - 1 = 0) \quad .2 = (4 [\cos(\theta)]^2 = 1) \\ .3 = (\tan(\theta) \cos(\theta) = \cos(\theta)) \quad .4 = (3 [\tan(\theta)]^2 - 1 = 0) \end{array} \right]
\end{aligned}$$

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00062XX

TrigonometryExercise6 for No.645129

$$\begin{aligned}
No1 &= \left[\begin{array}{l} .1 = (2 \sin(\theta) + 1 = 0) \quad .2 = (4 [\cos(\theta)]^2 - 1 = 0) \\ .3 = (\sin(2\theta) = 0) \quad .4 = (\cos(3\theta) = 1) \end{array} \right] \\
No2 &= \left[\begin{array}{l} .1 = (2 [\sin(\theta)]^2 - 3 \sin(\theta) + 1 = 0) \quad .2 = (2 [\cos(\theta)]^2 - \sin(\theta) - 1 = 0) \\ .3 = (2 [\sin(\theta)]^2 - \sin(\theta) = 0) \quad .4 = (\sin(\theta) - \cos(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \cos(\theta) = \cos(\theta)) \quad .6 = (4 [\sin(\theta)]^3 - \sin(\theta) = 0) \end{array} \right] \\
No3 &= \left[\begin{array}{l} .1 = (2 \cos(\theta) - 1 = 0) \quad .2 = (4 [\cos(\theta)]^2 - 3 = 0) \\ .3 = (\cos(4\theta) = 0) \quad .4 = (\sin(4\theta) = 1) \end{array} \right] \\
No4 &= \left[\begin{array}{l} .1 = (2 [\cos(\theta)]^2 + 3 \cos(\theta) + 1 = 0) \quad .2 = (2 [\cos(\theta)]^2 + \sin(\theta) - 1 = 0) \\ .3 = (2 [\sin(\theta)]^2 + \sin(\theta) = 0) \quad .4 = (\cos(\theta) - \sin(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \sin(\theta) = \tan(\theta)) \quad .6 = (4 [\cos(\theta)]^3 - \cos(\theta) = 0) \end{array} \right] \\
No5 &= \left[\begin{array}{l} .1 = (2 \sin(\theta) - 1 = 0) \quad .2 = (4 [\sin(\theta)]^2 = 1) \\ .3 = (\tan(\theta) \sin(\theta) = \sin(\theta)) \quad .4 = ([\tan(\theta)]^2 - 3 = 0) \end{array} \right]
\end{aligned}$$

X [Page = 0031] XXX

:
:
:
:
:
:
:
:
:
:
:

XX

:
:
:
:
:

X Math@MUT XXXM5/1-6400306-00063XX

TrigonometryExercise6 for No.645130

$$No1 = \left[\begin{array}{ll} .1 = (2 \sin(\theta) + 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 1 = 0) \\ .3 = (\sin(3 \theta) = 0) & .4 = (\sin(4 \theta) = 1) \end{array} \right]$$

$$No2 = \left[\begin{array}{ll} .1 = (2 [\sin(\theta)]^2 + 3 \sin(\theta) + 1 = 0) & .2 = (2 [\sin(\theta)]^2 + 3 \cos(\theta) - 3 = 0) \\ .3 = (2 [\cos(\theta)]^2 - \cos(\theta) = 0) & .4 = (\cos(\theta) - \sin(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \sin(\theta) = \tan(\theta)) & .6 = (4 [\sin(\theta)]^3 - \sin(\theta) = 0) \end{array} \right]$$

$$No3 = \left[\begin{array}{ll} .1 = (2 \sin(\theta) - 1 = 0) & .2 = (4 [\cos(\theta)]^2 - 3 = 0) \\ .3 = (\cos(4 \theta) = 0) & .4 = (\sin(2 \theta) = 1) \end{array} \right]$$

$$No4 = \left[\begin{array}{ll} .1 = (2 [\sin(\theta)]^2 + \sin(\theta) - 1 = 0) & .2 = (2 [\cos(\theta)]^2 + \sin(\theta) - 1 = 0) \\ .3 = (2 [\cos(\theta)]^2 + \cos(\theta) = 0) & .4 = (\sin(\theta) + \cos(\theta) = \sqrt{2}) \\ .5 = (\tan(\theta) \cos(\theta) = \cos(\theta)) & .6 = (4 [\cos(\theta)]^3 - \cos(\theta) = 0) \end{array} \right]$$

$$No5 = \left[\begin{array}{ll} .1 = (2 \cos(\theta) - 1 = 0) & .2 = (4 [\sin(\theta)]^2 = 1) \\ .3 = (\tan(\theta) \sin(\theta) = \sin(\theta)) & .4 = (3 [\tan(\theta)]^2 - 1 = 0) \end{array} \right]$$

XX

[>