

$$\begin{aligned}
 & \text{Ans2} = \left[\begin{array}{l}
 .1 = [\lim_{x \rightarrow 0^-} f(x) = -2, \lim_{x \rightarrow 0^+} f(x) = 2, \lim_{x \rightarrow 0} f(x) = \text{undefined}] \\
 .2 = [\lim_{x \rightarrow 0^-} g(x) = 3, \lim_{x \rightarrow 0^+} g(x) = 3, \lim_{x \rightarrow 0} g(x) = 3] \\
 .3 = \left[\begin{array}{lll}
 \lim_{x \rightarrow 0^-} f(x) + g(x) = 1 & \lim_{x \rightarrow 0^+} f(x) - g(x) = -1 & \lim_{x \rightarrow 0^-} f(x) g(x) = -6 \\
 \lim_{x \rightarrow 0^+} \frac{f(x)}{g(x)} = \frac{2}{3} & \lim_{x \rightarrow 0} g(x) - f(x) = \text{undefined} & \text{Math@MUT}
 \end{array} \right]
 \end{array} \right] \\
 & \text{Ans3} = \left[\begin{array}{lll}
 .1 = (\lim_{x \rightarrow 1^-} f(x) = 0) & .2 = (\lim_{x \rightarrow 1^+} f(x) = 1) & .3 = (\lim_{x \rightarrow 1} f(x) = \text{undefined}) \\
 .4 = (\lim_{x \rightarrow (-2.8)^-} f(x) = -3.) & .5 = (\lim_{x \rightarrow (-2.8)^+} f(x) = -3.) & .6 = (\lim_{x \rightarrow (-2.8)} f(x) = -3.)
 \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 \text{Ans2} &= \left[\begin{array}{l} .1 = [\lim_{x \rightarrow 0^-} f(x) = -3, \lim_{x \rightarrow 0^+} f(x) = 3, \lim_{x \rightarrow 0} f(x) = \text{undefined}] \\ .2 = [\lim_{x \rightarrow 0^-} g(x) = -3, \lim_{x \rightarrow 0^+} g(x) = -3, \lim_{x \rightarrow 0} g(x) = -3] \\ .3 = \left[\begin{array}{lll} \lim_{x \rightarrow 0^+} f(x) + g(x) = 0 & \lim_{x \rightarrow 0^-} f(x) - g(x) = 0 & \lim_{x \rightarrow 0^+} f(x) g(x) = -9 \\ \lim_{x \rightarrow 0^-} \frac{f(x)}{g(x)} = 1 & \lim_{x \rightarrow 0} \frac{f(x)}{g(x)} = \text{undefined} & \text{Math@MUT} \end{array} \right] \end{array} \right] \\
 \text{Ans3} &= \left[\begin{array}{lll} .1 = (\lim_{x \rightarrow 1^-} f(x) = 0) & .2 = (\lim_{x \rightarrow 1^+} f(x) = 1) & .3 = (\lim_{x \rightarrow 1} f(x) = \text{undefined}) \\ .4 = (\lim_{x \rightarrow (-3.2)^-} f(x) = -4.) & .5 = (\lim_{x \rightarrow (-3.2)^+} f(x) = -4.) & .6 = (\lim_{x \rightarrow (-3.2)} f(x) = -4.) \end{array} \right]
 \end{aligned}$$

[illegible]